

On the influence of Stark broadening on Cr I lines in stellar atmospheres

M. S. Dimitrijević^{1,2}, T. Ryabchikova^{3,4}, L. Č. Popović^{1,2,5}, D. Shulyak⁶, and S. Khan⁶

¹ Astronomical Observatory, Volgina 7, 11160 Belgrade 74, Serbia

² Isaac Newton Institute of Chile, Yugoslavia Branch

³ Institute of Astronomy, Russian Academy of Science, Pyatnitskaya 48, 119017 Moscow, Russia

⁴ Institute for Astronomy, University of Vienna, Türkenschanzstrasse 17, A-1180 Vienna, Austria

⁵ Astrophysikalisches Institut Potsdam, An der Sternwarte 16, 14482 Potsdam, Germany

⁶ Tavrian National University, Yaltinskaya 4, 330000 Simferopol, Crimea, Ukraine

Received / Accepted

Abstract. Using the semiclassical perturbation method, electron-, proton-, and ionized helium-impact line widths and shifts for the nine Cr I spectral lines from the $4p^7P^0 - 4d^7D$ multiplet, have been calculated for a perturbers density of 10^{14} cm^{-3} and temperatures $T = 2,500 - 50,000 \text{ K}$. The obtained results have been used to investigate the influence of Stark broadening effect in the Cr-rich Ap star β CrB atmosphere on line shapes of these lines. It has been found that the contribution of the proton and He II collisions to the line width and shift is significant, and it is comparable and sometimes (depending of the electron temperature) even larger than electron-impact contribution. Moreover, not only the Stark line width, but, depending on the electron-, proton-, and He II density in stellar atmosphere also the Stark shift may contribute to the blue as well as to the red asymmetry of the same line. The obtained results have been used to investigate the influence of Stark broadening effect on line shapes of Cr I lines in the atmosphere of Cr-rich Ap star β CrB.

Key words. atomic processes: Stark effect – line: profiles – stars: chemically peculiar – stars: individual: β CrB

1. Introduction

The Stark broadening is the most significant pressure broadening mechanism for A and B stars and this effect should be taken into account in investigation, analysis and modelling of their atmospheres. In our previous works (Popović et al. 1999, Popović et al. 2001, Dimitrijević et al. 2003) we have shown that the Stark broadening may change the spectral line equivalent widths by 10-45%, hence neglecting this mechanism, a significant errors in abundance determinations may be introduced. Stark data become extremely important after discovering abundance gradients in the atmospheres of magnetic peculiar (Ap) stars (Babel 1992, Ryabchikova et al. 2002, Wade et al. 2003). High resolution spectra allow us to perform stratification analysis using line profiles, and strong lines with developed wings provide us with the most accurate information about distribution of the element through the stellar atmosphere (see Dimitrijević et al. 2003 for Si).

Chromium is one of the most anomalous elements in Ap stars. It was shown to be concentrated in the deeper atmospheric layers in Ap stars β CrB (Wade et al. 2003) and

in γ Equ (Ryabchikova et al. 2002), where electron density is high enough to favour the Stark effect. The most of Cr I and Cr II lines in the optical spectral region have rather small Stark damping constants and hence, no measurable Stark wings appeared. However, Cr I lines from $4p - 4d$ transitions are known to have rather large Stark damping constants according to calculations made by Kurucz (1993).

We present new calculations of Cr I lines Stark widths and shifts based on the semiclassical perturbation approach (Sahal-Bréchet 1969a,b, 1974, Dimitrijević & Sahal-Bréchet 1984a). Method of these calculations is described in Section 2, magnetic synthetic spectrum calculations are described in Section 3. Results of Stark broadening calculations are given in Section 4.1. An application of new data to spectrum synthesis in magnetic Ap star β CrB is considered in Section 4.2.

2. The Stark broadening parameter calculation

Calculations have been performed within the semiclassical perturbation formalism, developed and discussed in detail in Sahal-Bréchet (1969ab). This formalism, as well as the corresponding computer code, have been

optimized and updated several times (see e.g. Sahal-Bréchet, 1974; Dimitrijević and Sahal-Bréchet, 1984a, Dimitrijević, 1996).

Within this formalism, the full width of a neutral emitter isolated spectral line broadened by electron impacts can be expressed in terms of cross sections for elastic and inelastic processes as

$$W_{if} = \frac{2\lambda_{if}^2}{2\pi c} n_e \int v f(v) dv \left(\sum_{i' \neq i} \sigma_{ii'}(v) + \sum_{f' \neq f} \sigma_{ff'}(v) + \sigma_{el} \right) \quad (1)$$

and the corresponding line shift as

$$d_{if} = \frac{\lambda_{if}^2}{2\pi c} n_e \int v f(v) dv \int_{R_3}^{R_D} 2\pi \rho d\rho \sin 2\phi_p \quad (2)$$

Here, λ_{if} is the wavelength of the line originating from the transition with the initial atomic energy level i and the final level f , c is the velocity of light, n_e is the electron density, $f(v)$ is the Maxwellian velocity distribution function for electrons, m is the electron mass, k is the Boltzmann constant, T is the temperature, and ρ denotes the impact parameter of the incoming electron. The inelastic cross section $\sigma_{jj'}(v)$ is determined according to Chapter 3 in Sahal-Bréchet (1969b), and elastic cross section σ_{el} according to Sahal-Bréchet (1969a). The cut-offs, included in order to maintain for the unitarity of the S -matrix, are described in Section 1 of Chapter 3 in Sahal-Bréchet (1969a).

The formulae for the ion-impact broadening parameters are analogous to the formulae for electron-impact broadening. For the colliding ions, the validity of the impact approximation has to be checked in the far wings.

3. Line profile calculations

Model atmosphere calculations as well as calculations of the absorption coefficients were made with the local thermodynamical equilibrium (LTE) approximation. Model calculations were performed with the ATLAS9 code (Kurucz 1993), modified by Piskunov & Kupka (2001) to include individual chemical composition of the star in line opacity calculations (see Kupka et al. 2004).

The next step is the calculation of the outward flux at corresponding wavelengths points using the given model. For this purpose we used SYNTHM code written by Khan (2004). This code allows to calculate synthetic spectra of early and intermediate type of stars taking into account magnetic field effects and stratification of chemical elements.

The computational scheme is as follows. For each line we find the central opacity as

$$\alpha_{line} = \frac{\pi e^2}{mc} g f_{if} e^{-\frac{\chi}{kT}} \frac{n}{\rho} (1 - e^{-\frac{h\nu}{kT}}), \quad (3)$$

where α_ν is the mass absorption coefficient at frequency ν , e is the electron charge, g is the statistical weight, f_{if} is

the oscillator strength for a given transition, χ is the excitation energy, n is the number density of a corresponding element in a given ionization stage multiplied by partition function, ρ is the density and h is the Planck constant. The last factor describes stimulated emission.

Next, we compute the total damping parameter

$$\gamma = \gamma_{rad} + \gamma_{Stark} + \gamma_{neutral}. \quad (4)$$

Here γ_{rad} , γ_{Stark} and $\gamma_{neutral}$ are the radiative broadening, Stark broadening and damping parameters due to neutral atom collisions respectively. The values of γ_{rad} , $\gamma_{neutral}$, excitation energy χ and oscillator strength gf were taken from the Vienna Atomic Line Database (VALD) (Kupka et al. 1999). In the case of neutral atom broadening we assumed that perturbing particles are atoms of neutral hydrogen and helium only (Van der Waals broadening). This assumption is applicable to almost all types of stars due to high hydrogen and helium cosmic abundances. We shall discuss the competition between the broadenings caused by the Stark effect and neutral hydrogen collisions in the atmospheres of our template star β CrB in Section 4.2.

In order to include Stark broadening effects we added the approximate formulas (see Eqs. (20) and (21) in Sec. 4.1) in the code. The Stark width and shift are

$$\gamma_{Stark} = \gamma_{Stark}^{(e)} n_e + \gamma_{Stark}^{(p)} n_p + \gamma_{Stark}^{(HeII)} n_{HeII}, \quad (5)$$

$$d_{Stark} = d_{Stark}^{(e)} n_e + d_{Stark}^{(p)} n_p + d_{Stark}^{(HeII)} n_{HeII}, \quad (6)$$

where n_e , n_p and n_{HeII} are the corresponding densities of electrons, protons and He II ions respectively. The resulting opacity profile is given by the Voigt function (Doppler + pressure broadening).

The next step is to solve transfer equation with new Stark damping parameters. In presence of magnetic field atomic level k determined by quantum numbers J_k , L_k , S_k splits into $2J_k + 1$ states with $M_k = -J_k, \dots, +J_k$. The absolute value of splitting is defined by field modulus $|\mathbf{B}|$ and Landé factor g_k which in the case of LS coupling is calculated as

$$g_k = \frac{3}{2} + \frac{S_k(S_k + 1) - L_k(L_k + 1)}{2J_k(J_k + 1)}. \quad (7)$$

The possible transitions between splitted levels (u - upper and l - lower level) are allowed by selection rules

$$\Delta M = M_u - M_l = \begin{cases} +1 \equiv b, \\ 0 \equiv p, \\ -1 \equiv r. \end{cases} \quad (8)$$

For a normal Zeeman triplet the subscript p corresponds to the unshifted π component, whereas b and r correspond to the blue- and red-shifted σ components respectively. In the general case of an anomalous splitting indices p , b , r refer to the series of the π and σ components.

The wavelength shift of the component relative to the laboratory line centre λ_0 is defined by

$$\Delta\lambda = \frac{e\lambda_0^2 |\mathbf{B}|}{4\pi mc^2} (g_l M_l - g_u M_u). \quad (9)$$

The relative strengths of components $S_{p,r,b}$ in accordance with Sobelman (1977) are proportional to

$$\begin{pmatrix} J_u & 1 & J_l \\ -M_u & M_u - M_l & M_l \end{pmatrix}^2, \quad (10)$$

where the last structure is a $3j$ -symbol. The normalization of components is done so that

$$\sum S_p = \sum S_b = \sum S_r = 1. \quad (11)$$

In general case the polarized radiation can be described by means of Stokes $IQUV$ parameters. The transfer equation is

$$\frac{d\mathbf{I}}{dz} = -\mathbf{K}\mathbf{I} + \mathbf{J}. \quad (12)$$

Here $\mathbf{I} = (I, Q, U, V)^T$ is the Stokes vector, $\mathbf{K}$ is the absorption matrix and $\mathbf{J}$ is the emission vector,

$$\mathbf{K} = \alpha_c \mathbf{1} + \sum_{lines} \alpha_{line} \Phi_{line}, \quad (13)$$

$$\mathbf{J}_\nu = \alpha_c B_\nu(T) \mathbf{e}_0 + B_\nu(T) \sum_{lines} \alpha_{line} \Phi_{line} \mathbf{e}_0, \quad (14)$$

where $\mathbf{1}$ is the 4×4 unit matrix, $\mathbf{e}_0 = (1, 0, 0, 0)^T$, α_c is the continuum absorption coefficient, α_{line} is the line central opacity for zero damping and zero magnetic field given by equation (3). Here we also assume LTE so the source function is equal to Planck function $B_\nu(T)$.

The line absorption matrix Φ_{line} is created using absorption profiles ϕ_j and anomalous dispersion profiles ψ_j of π and $\sigma_\pm$ components in accordance with Rees et al. (1989). These profiles are calculated as (for $j = p, b, r$)

$$\phi_j = \sum_{i_j=1}^{N_j} S_{i_j} H\left(a, v - \frac{\Delta\lambda_{i_j}}{\Delta\lambda_D}\right), \quad (15)$$

$$\psi_j = 2 \sum_{i_j=1}^{N_j} S_{i_j} F\left(a, v - \frac{\Delta\lambda_{i_j}}{\Delta\lambda_D}\right), \quad (16)$$

where $H(a, u)$ and $F(a, u)$ are the Voigt and Faraday-Voigt functions,

$$a = \frac{\gamma \lambda_0^2}{4\pi c \Delta\lambda_D}, \quad (17)$$

$$v = \frac{\lambda - \lambda_0 + d_{Stark}}{\Delta\lambda_D}. \quad (18)$$

The Stark shift d_{Stark} and the damping parameter γ have been found from (6) and (4), respectively. The Doppler width $\Delta\lambda_D$ is

$$\Delta\lambda_D = \frac{\lambda}{c} \sqrt{\frac{2kT}{m_A} + \xi_t^2}, \quad (19)$$

where m_A is the mass of the absorber atom and ξ_t is the microturbulent velocity.

The transfer equation (12) is solved by DELO method (Rees et al. 1989) with quadratic approximation for the source function which was suggested by Socas-Navarro et al. (2000) and written in Fortran by Piskunov & Kochukhov (2000). Also, we used fast Humlicek (1982) algorithm for approximations of Voigt and Faraday-Voigt functions.

The convolution of the synthetic spectra due to rotation of the star was produced by IDL procedure of Valenti & Anderson implemented in the ROTATE IDL code by N. Piskunov.

4. Results

4.1. Stark broadening data

The atomic energy levels needed for Stark broadening calculations were taken from Wiese and Musgrove (1989). Oscillator strengths have been calculated by using the method of Bates and Damgaard (1949) and tables of Oertel and Shomo (1968). For higher levels, the method described in van Regemorter, Binh Dy and Prud'homme (1979) has been applied.

The spectrum of neutral chromium is complex and not known well enough for a good calculation of the considered lines. The principal problem is the absence of the reliable experimental data for f levels. However, by inspecting theoretically predicted Cr I levels in Moore (1971), one can see that the $4f$ level is missing. In accordance with the decrease of distance between existing s, p , and d levels, we estimated that the eventual $5f$ level should be around 52000 cm^{-1} and not closer than 50000 cm^{-1} . We checked results without $5f$ level and with a fictive $5f$ level at 52000 and 50000 cm^{-1} . In all cases, line widths differed by less than 1%, while the shift varies within the limits of several percents. Consequently we performed calculations without the contribution of f energy levels. Since the average estimated error of the semiclassical method is $\pm 30\%$, due to additional approximations and uncertainties, we estimate the error bars of our results to be $\pm 50\%$.

Our results for electron-, proton-, and ionized helium-impact line widths and shifts for the nine considered Cr I spectral lines for a perturbed density of 10^{14} cm^{-3} and temperatures $T = 2,500 - 50,000 \text{ K}$, are shown in Table 1. For perturber densities lower than those tabulated here, Stark broadening parameters vary linearly with perturber density. The nonlinear behavior of Stark broadening parameters at higher densities is the consequence of the influence of Debye shielding and has been analyzed in detail in Dimitrijević and Sahal-Bréchet (1984b).

Table 1. Stark broadening parameters for Cr I $4p - 4d$ spectral lines. This table shows electron-, proton-, and ionized helium-impact broadening parameters for Cr I for a perturber density of 10^{14} cm^{-3} and temperatures from 2,500 up to 50,000 K. The quantity C (given in $\text{\AA cm}^{-3}$), when divided by the corresponding full width at half maximum, gives an estimate for the maximum perturber density for which tabulated data may be used. For higher densities, the isolated line approximation used in the calculations breaks down. The asterisk identifies cases for which the collision volume multiplied by the perturber density (the condition for validity of the impact approximation) lies between 0.1 and 0.5. When this value is larger than 0.5, the corresponding results have been omitted. $\text{FWHM}(\text{\AA})$ denotes full line width at half maximum in $\text{\AA}$, while $\text{SHIFT}(\text{\AA})$ denotes line shift in $\text{\AA}$.

PERTURBERS ARE:		ELECTRONS		PROTONS		HELIUM IONS	
TRANSITION	T(K)	FWHM(A)	SHIFT(A)	FWHM(A)	SHIFT(A)	FWHM(A)	SHIFT(A)
Cr I $^4P_2^o - ^4D_1$ 5276.07 A C = 0.43E+15	2500.	0.890E-02	-0.205E-02	0.461E-02	-0.379E-02	0.344E-02	-0.299E-02
	5000.	0.772E-02	-0.146E-02	0.551E-02	-0.437E-02	0.388E-02	-0.346E-02
	10000.	0.655E-02	-0.104E-02	0.678E-02	-0.505E-02	0.439E-02	-0.396E-02
	20000.	0.548E-02	-0.767E-03	0.832E-02	-0.592E-02	0.503E-02	-0.451E-02
	30000.	0.493E-02	-0.660E-03	0.915E-02	-0.651E-02	0.547E-02	-0.488E-02
Cr I $^4P_2^o - ^4D_2$ 5275.75 A C = 0.11E+14	2500.	0.193E-01	-0.216E-02				
	5000.	0.164E-01	-0.155E-02				
	10000.	0.136E-01	-0.108E-02				
	20000.	0.111E-01	-0.790E-03	*0.717E-01	-0.298E-01		
	30000.	0.983E-02	-0.675E-03	*0.702E-01	-0.273E-01		
Cr I $^4P_2^o - ^4D_3$ 5275.28 A C = 0.60E+14	2500.	0.240E-01	-0.939E-03	*0.216E-01	-0.138E-01	*0.131E-01	-0.102E-01
	5000.	0.210E-01	-0.677E-03	*0.253E-01	-0.170E-01	*0.151E-01	-0.124E-01
	10000.	0.178E-01	-0.473E-03	*0.274E-01	-0.193E-01	*0.169E-01	-0.150E-01
	20000.	0.147E-01	-0.401E-03	0.280E-01	-0.193E-01	*0.170E-01	-0.179E-01
	30000.	0.131E-01	-0.393E-03	0.283E-01	-0.180E-01	*0.157E-01	-0.193E-01
Cr I $^4P_3^o - ^4D_2$ 5298.49 A C = 0.12E+14	2500.	0.195E-01	-0.219E-02				
	5000.	0.165E-01	-0.156E-02				
	10000.	0.137E-01	-0.109E-02				
	20000.	0.112E-01	-0.801E-03	*0.723E-01	-0.301E-01		
	30000.	0.992E-02	-0.685E-03	*0.708E-01	-0.276E-01		
Cr I $^4P_3^o - ^4D_3$ 5298.02 A C = 0.60E+14	2500.	0.243E-01	-0.950E-03	*0.218E-01	-0.140E-01	*0.133E-01	-0.103E-01
	5000.	0.212E-01	-0.687E-03	*0.256E-01	-0.171E-01	*0.153E-01	-0.125E-01
	10000.	0.179E-01	-0.481E-03	*0.276E-01	-0.195E-01	*0.171E-01	-0.151E-01
	20000.	0.148E-01	-0.408E-03	0.283E-01	-0.195E-01	*0.171E-01	-0.180E-01
	30000.	0.132E-01	-0.400E-03	0.286E-01	-0.182E-01	*0.158E-01	-0.195E-01
Cr I $^4P_3^o - ^4D_3$ 5298.02 A C = 0.60E+14	50000.	0.113E-01	-0.400E-03	0.291E-01	-0.157E-01	*0.128E-01	-0.201E-01

After testing the density dependence of Stark parameters, we have found that the width and shift are linear functions of density for perturber densities smaller than 10^{16} cm^{-3} and can be scaled by the simple formula:

$$(W, d)_N = (W, d)_0 \left(\frac{N}{10^{14}} \right), \quad (20)$$

where $(W, d)_N$ are the width and shift at a perturber density $N \text{ (cm}^{-3}\text{)}$, and $(W, d)_0$ are width and shift given in Table 1, respectively.

In order to simplify the use of Stark broadening data in the codes for stellar spectral synthesis, we have found an analytical expression for Stark widths and shifts


$$\frac{W}{n_e} [\text{\AA}] = c_1 \cdot (A + T^B), \quad (21)$$

$$\frac{d}{n_e} [\text{\AA}] = c_2 \cdot (A + C \cdot T^B). \quad (22)$$

The constants c_1 , c_2 , A , B and C are given in Table 2. We take T as $T/(10000\text{K})$. The analytical fit is present in Figs. 1 and 2. As one can see from the Figs. 1 and 2, the analytical assumption good fit the calculated value in both, Stark widths and shifts. Here we should note that in some lines (as e.g. in Cr I $\lambda = 5329.14 \text{ \AA}$) the analytical fit in the case of proton and He II impact broadening parameters not exactly fit the calculated values (see e.g. Fig. 2, fit for proton-impact shift), but the differences are smaller than the expected error of our calculations ($\approx \pm 50\%$).


Table 1. (continued)

PERTURBERS ARE:		ELECTRONS		PROTONS		HELIUM IONS	
TRANSITION	T(K)	FWHM(A)	SHIFT(A)	FWHM(A)	SHIFT(A)	FWHM(A)	SHIFT(A)
Cr I $^7P_3^o - ^7D_4$ 5297.38 Å C = 0.12E+15	2500.	0.182E-01	0.452E-03	0.109E-01	-0.737E-02	*0.683E-02	-0.564E-02
	5000.	0.156E-01	0.277E-03	0.130E-01	-0.902E-02	*0.788E-02	-0.664E-02
	10000.	0.131E-01	0.129E-03	0.144E-01	-0.102E-01	0.909E-02	-0.787E-02
	20000.	0.108E-01	-0.434E-04	0.168E-01	-0.945E-02	0.956E-02	-0.934E-02
	30000.	0.956E-02	-0.141E-03	0.172E-01	-0.830E-02	0.945E-02	-0.100E-01
	50000.	0.822E-02	-0.240E-03	0.194E-01	-0.603E-02	0.793E-02	-0.100E-01
Cr I $^7P_4^o - ^7D_3$ 5329.78 Å C = 0.61E+14	2500.	0.245E-01	-0.956E-03	*0.220E-01	-0.141E-01	*0.134E-01	-0.104E-01
	5000.	0.214E-01	-0.701E-03	*0.259E-01	-0.173E-01	*0.155E-01	-0.127E-01
	10000.	0.181E-01	-0.492E-03	*0.280E-01	-0.197E-01	*0.173E-01	-0.153E-01
	20000.	0.150E-01	-0.419E-03	0.286E-01	-0.197E-01	*0.173E-01	-0.182E-01
	30000.	0.134E-01	-0.411E-03	0.289E-01	-0.184E-01	*0.160E-01	-0.197E-01
	50000.	0.115E-01	-0.410E-03	0.295E-01	-0.159E-01	*0.130E-01	-0.204E-01
Cr I $^7P_4^o - ^7D_4$ 5329.14 Å C = 0.12E+15	2500.	0.184E-01	0.460E-03	0.110E-01	-0.746E-02	*0.692E-02	-0.570E-02
	5000.	0.158E-01	0.276E-03	0.132E-01	-0.911E-02	*0.798E-02	-0.672E-02
	10000.	0.132E-01	0.127E-03	0.146E-01	-0.103E-01	0.920E-02	-0.796E-02
	20000.	0.109E-01	-0.484E-04	0.170E-01	-0.956E-02	0.967E-02	-0.946E-02
	30000.	0.968E-02	-0.147E-03	0.174E-01	-0.840E-02	0.956E-02	-0.101E-01
	50000.	0.832E-02	-0.248E-03	0.197E-01	-0.610E-02	0.802E-02	-0.101E-01
Cr I $^7P_4^o - ^7D_5$ 5328.32 Å C = 0.40E+15	2500.	0.920E-02	0.162E-02	0.499E-02	0.407E-02	0.370E-02	0.321E-02
	5000.	0.800E-02	0.979E-03	0.599E-02	0.470E-02	0.417E-02	0.372E-02
	10000.	0.678E-02	0.474E-03	0.739E-02	0.544E-02	0.473E-02	0.426E-02
	20000.	0.567E-02	0.103E-03	0.902E-02	0.641E-02	0.543E-02	0.485E-02
	30000.	0.509E-02	-0.604E-04	0.987E-02	0.709E-02	0.590E-02	0.526E-02
	50000.	0.445E-02	-0.208E-03	0.106E-01	0.784E-02	0.647E-02	0.589E-02

**Fig. 1.** The analytic fit of Cr I $\lambda = 5329.14$ Å (upper Fig.) and $\lambda = 5328.32$ Å (down) Stark widths due to impact with: electrons (open circles), protons (full circles) and He II (full triangles).

4.2. The Stark broadening effect on the shape of Cr I lines

In spite of the rather large Stark damping constants the effect is not observable in stars with solar Cr abundance. In hot stars where electron and proton densities are high, Cr I

**Fig. 2.** The same as in Fig. 1, but for Stark shift.

lines considered are generally very weak, while in cooler stars (solar type) where these lines are strong enough other broadening effects are more significant. The only chance to look at Stark effect is the stratified atmosphere of Cr-rich Ap star. The best example is the well known magnetic star β CrB.


Our spectral analysis was based on the spectrum of β CrB obtained in February 1998 with MuSiCoS spectropolarimeter mounted on the 2 m telescope at Pic du

Table 2. The parameters A, B and C of the approximate formulae for Stark widths and shifts.

Line	5276.07 Å	5275.75 Å	5275.28 Å	5298.49 Å	5298.02 Å	5297.38 Å	5329.78 Å	5329.14 Å	5328.32 Å
WIDTH									
electrons									
c_1	1E-16	1E-15	1E-15	1E-15	1E-15	1E-15	1E-15	1E-15	1E-16
A	-0.3474	-0.86156	-0.8218	-0.86039	-0.82019	-0.8674	-0.81818	-0.86596	-0.325497
B	-0.15596	-0.03684	-0.04346	-0.03716	-0.04412	-0.0337	-0.04412	-0.03410	-0.161576
protons									
c_1	1E-16	1E-15	1E-15	1E-15	1E-15	1E-15	1E-15	1E-15	1E-15
A	-0.3194	-0.23723*	-0.7393	-0.2312*	-0.7368	-0.85357	-0.7338	-0.85169	-0.924767
B	0.1789	-0.06276*	0.02231	-0.062768*	0.02238	0.02723	0.02281	0.027692	0.01959
He II									
c_1	1E-16	-	1E-15	-	1E-15	1E-16	1E-15	1E-16	1E-16
A	-0.5525	-	-0.84377	-	-0.84219	-0.14236	-0.84037*	-0.13199	-0.51847
B	0.08565	-	0.01192	-	0.011451	0.11010	0.01177*	0.110894	0.092283
SHIFT									
electrons									
c_2	1E-16	1E-16	1E-17	1E-16	1E-17	1E-17	1E-17	1E-17	1E-16
A	-3.4929	-6.7617	-81.5107	-2.5259	-9.0041	49.6736	-16.5511	40.5533	23.3420
B	0.01451	0.00793	0.00221	0.02206	0.02144	0.00526	0.01131	0.00586	0.00261
C	3.3728	6.6362	80.9250	2.3985	8.4079	-49.60704	15.9454	-40.4305	-23.2800
protons									
c_2	1E-16	1E-15	1E-15	1E-15	1E-15	1E-16	1E-15	1E-16	1E-16
A	-0.0170	0.25969*	3.0908	-3.3759*	0.5500	-2.4538	3.7646	18.8373	50.1882
B	0.22391	-0.122027*	0.00215	0.00846*	0.00990	0.02228	0.00188	-0.00173	-0.002557
C	-0.4916	-0.6076*	-3.2607	3.0621*	-0.7217	1.6060	-3.9381	-19.6934	-49.62339
He II									
c_2	1E-16	-	1E-15	-	1E-15	1E-16	1E-15	1E-16	1E-16
A	-0.0174	-	1.7973	-	2.4949	37.4669	2.0440	36.5920	0.0405
B	0.20567	-	0.01778	-	0.01314	0.00418	0.01604	0.00431	0.22021
C	-0.3771	-	-1.9474	-	-2.6462	-38.2561	-2.1971	-37.3904	0.3832

Midi observatory ($R=35000$). This spectrum was used by Wade et al. (2001,2003) in stratification analysis. Starting Cr and Fe distributions were taken from this analysis and were slightly corrected using a set of spectral lines of Cr I Cr II Fe I Fe II with different excitation energies of the lower level. The adopted distributions are shown in Fig. 3. Fe stratification is necessary to account correctly for blends.

Synthetic spectrum calculations in the regions of Cr I lines considered in the present paper was performed following the method described in Section 3. The surface magnetic field $B_s=5.4$ kG was derived from the magnetically splitted lines and used in all calculations. In the process of calculations we found that Cr I lines of our interest have incorrect wavelengths with the shifts up to 0.05 Å. We adjusted wavelengths calculating solar synthetic spectrum and comparing it with the Solar Flux Atlas (Kurucz et al. 1984). Improved wavelengths of Cr I lines derived by this way agree within 0.005 Å and better with the precise measurements by J. E. Murray (1992), kindly provided to us by R. Kurucz¹. We placed Murray's wavelengths in Table 1 and 2 and used them in spectral synthesis. We also estimated the contribution of Van der Waals broadening

**Fig. 3.** Empirically derived Cr and Fe distributions in the atmosphere of magnetic Ap star β CrB.

and found it to be by the order of magnitude smaller than Stark broadening through the whole atmosphere above $\log(\tau_{5000}) = -0.2$ with the rapid decrease below the photosphere.

We varied Stark broadening parameters given in Table 1 to achieve the best fit of the calculated Cr I line

¹ <http://cfaku5.cfa.harvard.edu/atoms/2400/>




Fig. 4. A comparison between synthetic spectrum calculations and the observed spectrum (thick line) of magnetic Ap star β CrB in the regions of Cr I lines 5297 Å (top panel), 5276 Å (middle panel) and 5329 Å (bottom panel). Full thin line - calculations with stratification shown in Fig 3 and Stark broadening data from Table 1 decreased by 70 %; dashed line - the same Stark broadening data but homogeneous Cr and Fe abundances (see text); dotted line - stratified Cr and Fe abundances, but Stark broadening calculated with the approximation formula.

profiles to the observations. Fig. 4 shows a comparison between observations and calculations in the region of 3 groups of Cr I lines. For comparison we present syn-

thetic calculations with homogeneous mean abundances $\log(Cr/N_{tot}) = -4.80$ and $\log(Fe/N_{tot}) = -4.15$ which were used in model atmosphere calculations (Kupka et al. 2004). To fit Cr I line wings we need to decrease current Stark widths by 60-70 %. The same order of overestimate of Stark damping constants, calculated within the semiclassical perturbation formalism, was obtained for Si II lines (see Dimitrijević et al. 2003). Any change in Cr distribution, which may be responsible for line profiles, too, is critical for nearby Cr I λ 5296.691 Å, which is insensitive to Stark broadening mechanism due to small value of Stark damping constant and is sensitive only to Cr distribution in stellar atmosphere. We also made a comparison with synthetic spectrum calculated using stratified Cr distribution and an approximation formula for Stark broadening (Cowley 1971). This formula is used in the cases when Stark broadening data, experimental or theoretical, are not available. Evidently, Stark broadening is overestimated when calculations are performed with this approximation formula.

For detailed investigation of the influence of Stark shifts on the observed line profile we calculated theoretical spectra with and without shifts for two separate Cr I lines 5297.377 Å and 5298.494 Å in homogeneous and stratified atmosphere of the non-rotating star. These two lines have the smallest and largest shift values, therefore one can make a relative comparison. Fig. 5 illustrates an influence of the Stark shift on line shape. The total shift in line position due to the Stark broadening effect is negligible for Cr I 5297.377 Å line both in homogeneous and stratified atmosphere. For the second line Stark broadening leads to theoretically measurable shift in the line position, in particular, in the case of Cr stratification resulting in a small line asymmetry. Unfortunately, in spectra of real stars effect is difficult to measure due to other broadening mechanisms (rotation, magnetic field) and mainly due to severe blending in the regions with Cr I lines from $4p - 4d$ transitions.

5. Conclusions

Stark broadening parameters for Cr I spectral lines from $4p^7P^0 - 4d^7D$ multiplet have been calculated and the influence of Stark broadening effect in stellar atmosphere for these lines, has been investigated. From our investigation we can conclude:

- (i) Although they belong to a same multiplet, the widths and shifts of the different lines can be quite different.
- (ii) The contribution of the proton and He II collisions to the line width and shift is significant, and it is comparable and sometimes (depending of the electron temperature) even larger than electron-impact contribution.
- (iii) Depending on the electron-, proton-, and He II density in stellar atmosphere the Stark shift may contribute to the blue as well as to the red asymmetry of the same line (see Fig. 2).




Fig. 5. Influence of Stark shift on Cr I line shapes in homogeneous (upper panel) and in stratified (lower panel) atmosphere. Calculations with both Stark widths and shifts are shown by solid lines while those without Stark shifts are shown by dashed line.

(iv) To fit well Cr I line wings we need to decrease the calculated Stark widths by 60-70%, which is the same order of overestimation as for Si I lines (Dimitrijević et al. 2003). The approximation formula of Cowley (1971), used in the cases where the adequate semiclassical calculation is not possible due to the lack of reliable atomic data, predicts also overestimated influence of Stark broadening in comparison with observations.

Acknowledgements. We are very thankful to G. Wade and R. Kurucz who provided us with the data necessary for our analysis. This work is a part of the projects GA-1195 “Influence of collisional processes on astrophysical plasma lineshapes” and GA-1196 “Astrophysical Spectroscopy of Extragalactic Objects” supported by the Ministry of Science and Environment protection of Serbia. The research was supported also by the Fonds zur Förderung der wissenschaftlichen Forschung P14984. TR thanks Leading Scientific School grant 162.2003.02 and the RFBR (grant 03-02-16342) for partial funding. L. Č. P. is supported by Alexander von Humboldt Foundation through the program for foreign scholars. DS acknowledges a financial support from INTAS grant 03-55-652.

References

Babel, J. 1992, *A&A*, 258, 449

- Bates, D. R., & Damgaard, A. 1949, *Trans.Roy.Soc. London*, Ser. A, 242, 101
- Cowley, C.R. 1971, *Obs.*, 91, 139
- Dimitrijević, M. S. 1996, *Zh. Priklad. Spektrosk.*, 63, 810
- Dimitrijević, M. S., & Sahal-Bréchet, S. 1984a, *JQSRT*, 31, 301
- Dimitrijević, M. S., & Sahal-Bréchet, S. 1984b, *A&A*, 136, 289
- Dimitrijević, M. S., Ryabchikova, T., Popović, L. Č., Shulyak, D., Tsymbal, V. 2003, *A&A*, 404, 1099
- Humlicek, J. 1982, *JQSRT*, 27, 437
- Khan, S. 2004, *JQSRT*, 88, 71
- Kupka, F., Paunzen, E., Iliev, I.Kh., & Maitzen, H.M. 2004, *MNRAS* 352, 863
- Kupka, F., Piskunov, N.E., Ryabchikova, T.A., Stempels, H.C., & Weiss, W.W. 1999, *A&AS*, 138, 119
- Kurucz, R.L. 1993, *CDROM13*, SAO, Cambridge
- Kurucz, R.L., Furenlid, I., Brault, J., & Testerman L. 1984, *NSO Atlas No. 1: Solar Flux Atlas from 296 to 1300 nm*, Sunspot, NSO
- Moore, C. E. 1971, *Atomic Energy Levels Vol. II*, NSRDS-NBS 35, U.S. Gov. Print. Office, Washington
- Murray, J. E. 1992, PhD thesis, Imperial College, London
- Oertel, G. K., & Shomo, L. P. 1968, *ApJS*, 16, 175
- Piskunov, N., & Kupka, F. 2001, *ApJ*, 547, 1040
- Piskunov, N., & Kochukhov, O. 2002, *A&A*, 381, 736
- Popović, L.C., Dimitrijević M. & Ryabchikova, T. 1999 *A&A*, 350, 719
- Popović, L.C., Simić S., Milovanović N., & Dimitrijević M. 2001, *ApJS*, 135, 109.
- Rees, D. E., Murphy, G. A., & Durrant, C. J. 1989, *ApJ*, 339, 1093
- Ryabchikova, T., Piskunov, N., Kochukhov, O., et al. 2002, *A&A*, 384, 545
- Ryabchikova, T., Wade, G., & LeBlanc, F. 2003, in *IAU Symposium No. 210, Modelling of Stellar Atmospheres*, eds. N.E. Piskunov, W.W. Weiss, & D.F. Gray, ASP, p. 301
- Sahal – Bréchet, S. 1969a, *A&A*, 1, 91
- Sahal – Bréchet, S. 1969b, *A&A*, 2, 322
- Sahal – Bréchet, S. 1974, *A&A*, 35, 321
- Sobelman, I. I. 1977, *Introduction to the theory of atomic spectra*, Nauka, Moscow
- Socas-Navarro, H., Trujillo Bueno, J., & Ruiz Cobo, B. 2000, *ApJ* 530, 97
- van Regemorter, H., Hoang Binh Dy, & Prud’homme, M. 1979, *J. Phys. B*, 12, 1073
- Wade, G. A., Ryabchikova, T. A., Bagnulo, S., & Piskunov, N. 2001, in *Magnetic Fields Across the Hertzsprung-Russell Diagram*, ed. G. Mathys, S. K. Solanki, & D. T. Wickramasinghe, ASP Conf. Ser., 248, 341
- Wade, G.A., LeBlanc, F., Ryabchikova, T.A., & Kudryavtsev, D. 2003, in *IAU Symp. 210 Modelling of Stellar Atmospheres*, eds. N.E. Piskunov, W.W. Weiss, & D.F. Gray, CD-D7
- Wiese, W.L., Musgrove, A. 1989, *Atomic Data for Fusion Vol. 6, Spectroscopic Data for Titanium, Chromium and Nickel*, Vol. 2. Chromium, Controlled Fusion Atomic Data Center, Oak Ridge National Laboratory, Oak Ridge